

AFMR in noncollinear antiferromagnet $\text{Mn}_3\text{Al}_2\text{Ge}_3\text{O}_{12}$.

Yulia Krasnikova^{1,2}, Vasiliy Glazkov^{1,2}, Timofey Soldatov³

¹P. L. Kapitza Institute for Physical Problems RAS, Moscow, Russia

² National Research University Higher School of Economics, Moscow, Russia

³ Moscow Institute of Physics and Technology, Dolgoprudny, Russia

Various magnetic garnet are actively studying for decades. The most popular is YIG (yttrium iron oxide), which has applications in UHF techniques [1], other garnets are used for magnetic cooling [2]. Unusual magnetic properties are partially due to frustration of exchange bonds. Garnet $\text{Mn}_3\text{Al}_2\text{Ge}_3\text{O}_{12}$ orders antiferromagnetically at $T_N=6.5$ K into complicated 12 sublattices noncollinear structure [3].

We have studied low energy dynamics of this ordered structure by means of antiferromagnetic resonance (AFMR). Experiments were performed at $T=1.8$ K, $f=1-120$ GHz, $H=0-6$ T. Representative $f(H)$ diagram for $H||[100]$ is shown in Figure 1. Experimental data were described by Andreev-Marchenko theory [4] using numerical modelling [5]. Additionally we have detected hysteresis of AFMR absorption connected with antiferromagnetic domains (Figure 2.). More detailed analysis of AFMR in $\text{Mn}_3\text{Al}_2\text{Ge}_3\text{O}_{12}$ is reported in [6].




Figure 1. AFMR $f(H)$ dependence at $H||[100]$, $T=1.8$ K. Solid lines - numeric calculations.




Figure 2. Hysteresis of AFMR signal. Arrows - field sweep direction. $H||[111]$, $f=37.6$ GHz, $T=1.8$ K. H^* - field of monodomenization.

- [1] E. J. J. Mallmann et al. Solid State Phenomena **202**, 65 (2013).
- [2] T. J. Sato Review of Scientific Instruments **87**, 123905 (2016).
- [3] A. Gukasov et al. J. Phys.: Condens. Matter **11**, 2869 (1999).
- [4] A. F. Andreev, V. I. Marchenko, UFN **130**, 1 (1980).
- [5] V. Glazkov et al., Appl. Magn. Reson. **47**, 1069 (2016).
- [6] Y. V. Krasnikova et al. Journal of Experimental and Theoretical Physics **125**, 3, 476 (2017) arXiv:1708.03961.